

ON THE CURVATURE OF SYMMETRIC PRODUCTS OF A COMPACT RIEMANN SURFACE

INDRANIL BISWAS

ABSTRACT. Let X be a compact connected Riemann surface of genus at least two. The main theorem of [BR] says that for any positive integer $n \leq 2(\text{genus}(X) - 1)$, the symmetric product $S^n(X)$ does not admit any Kähler metric satisfying the condition that all the holomorphic bisectional curvatures are nonnegative. Our aim here is to give a very simple and direct proof of this result of Bökstedt and Romão.

1. INTRODUCTION

The main theorem of [BR] says the following (see [BR, Theorem 1.1]):

Let X be a compact connected Riemann surface of genus at least two. If

$$1 \leq n \leq 2(\text{genus}(X) - 1),$$

then the symmetric product $S^n(X)$ does not admit any Kähler metric for which all the holomorphic bisectional curvatures are nonnegative.

Our aim here is to give a simple proof of this theorem.

In [Bi], the following related was proved (see [Bi, Theorem 1.1]):

Let X be a compact connected Riemann surface of genus at least two. If

$$n > 2(\text{genus}(X) - 1),$$

then $S^n(X)$ does not admit any Kähler metric for which all the holomorphic bisectional curvatures are nonnegative.

2. NONNEGATIVE HOLOMORPHIC BISECTIONAL CURVATURE

Let X be a compact connected Riemann surface of genus g , with $g \geq 2$. For any positive integer n , the n -fold symmetric product of X will be denoted by $S^n(X)$. We recall that $S^n(X)$ is the quotient of X^n by the natural action of the group of permutations P_n of the index set $\{1, \dots, n\}$. Let

$$(2.1) \quad q : X^n \longrightarrow S^n(X) := X^n/P_n$$

be the quotient map.

Theorem 2.1. *Take any integer $n \in [1, 2(g - 1)]$. The symmetric product $S^n(X)$ does not admit any Kähler metric satisfying the condition that all the holomorphic bisectional curvatures are nonnegative.*

2000 *Mathematics Subject Classification.* 14C20, 32Q10.

Key words and phrases. Riemann surface, symmetric product, holomorphic bisectional curvature.

Proof. Let $K_{X^n}^{-1} = \bigwedge^n TX^n$ and $K_{S^n(X)}^{-1} = \bigwedge^n TS^n(X)$ be the anti-canonical line bundles of X^n and $S^n(X)$ respectively. Fix distinct $n-1$ points $\{x_1, \dots, x_{n-1}\}$ of X . Let

$$(2.2) \quad \varphi : X \longrightarrow X^n$$

be the embedding defined by $x \mapsto \{x, x_1, \dots, x_{n-1}\} \in X^n$. Define

$$(2.3) \quad \tilde{\varphi} := q \circ \varphi : X \longrightarrow S^n(X),$$

where q is defined in (2.1).

We will show that

$$(2.4) \quad \text{degree}(\tilde{\varphi}^* K_{S^n(X)}^{-1}) = n - 2g + 1.$$

For $1 \leq i < j \leq n$, let $D_{i,j} \subset X^n$ be the divisor consisting of all $\{y_1, \dots, y_n\}$ such that $y_i = y_j$. Let

$$D := \sum_{1 \leq i < j \leq n} D_{i,j}$$

be the divisor on X^n . For the map q in (2.1), we have

$$q^* K_{S^n(X)}^{-1} = K_{X^n}^{-1} \otimes \mathcal{O}_{X^n}(D).$$

Therefore,

$$(2.5) \quad \tilde{\varphi}^* K_{S^n(X)}^{-1} = \varphi^* q^* K_{S^n(X)}^{-1} = (\varphi^* K_{X^n}^{-1}) \otimes (\varphi^* \mathcal{O}_{X^n}(D)),$$

where φ is the map in (2.2).

Now,

$$\varphi^* K_{X^n}^{-1} = TX \quad \text{and} \quad \varphi^* \mathcal{O}_{X^n}(D) = \mathcal{O}_X\left(\sum_{i=1}^{n-1} x_i\right),$$

where TX is the holomorphic tangent bundle of X . Therefore, from (2.5) we conclude that

$$\tilde{\varphi}^* K_{S^n(X)}^{-1} = (TX) \otimes \mathcal{O}_X\left(\sum_{i=1}^{n-1} x_i\right).$$

Hence

$$\text{degree}(\tilde{\varphi}^* K_{S^n(X)}^{-1}) = 2 - 2g + n - 1 = n - 2g + 1,$$

and (2.4) is proved.

Since $n \leq 2g - 2$, from (2.4) we conclude that

$$\text{degree}(\tilde{\varphi}^* K_{S^n(X)}^{-1}) < 0.$$

This immediately implies that $S^n(X)$ does not admit a Kähler metric for which all the holomorphic bisectional curvatures are nonnegative. \square

The Chern class $c_1(K_{S^n(X)}^{-1})$ is computed in [Ma, p. 333] and [ACGH, p. 323]. It should be possible to derive (2.4) using this description of $c_1(K_{S^n(X)}^{-1})$.

REFERENCES

- [ACGH] E. Arbarello, M. Cornalba, P. A. Griffiths and J. Harris, *Geometry of algebraic curves*. Vol. I, Grundlehren der Mathematischen Wissenschaften, 267, Springer–Verlag, New York, 1985.
- [Bi] I. Biswas, On Kähler structures over symmetric products of a Riemann surface, *Proc. of the A.M.S.*, <http://arxiv.org/abs/1106.3733>.
- [BR] M. Bökstedt and N. M. Romão, On the curvature of vortex moduli spaces, <http://arxiv.org/abs/1010.1488>.
- [Ma] I. G. Macdonald, Symmetric products of an algebraic curve, *Topology* **1** (1962), 319–343.

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD,
BOMBAY 400005, INDIA

E-mail address: `indranil@math.tifr.res.in`